


DUAL FUNCTIONING EXCIPIENT FOR METAL
CHELATE CONTRAST AGENTS

5 This is a continuation-in-part application
of U. S. Ser. No. 514,468 filed April 25, 1990.

10 The use of metal chelates as contrast agents
for magnetic resonance, x-ray, ultrasound, and
radiodiagnostic imaging is well known. Metal
chelates of the transition metal and lanthanide
metal ions are of primary interest due to their
strong paramagnetic and x-ray absorbing properties
as well as others. Because the free metal ions are
in general more toxic, these type of
15 pharmaceuticals are prepared in the form of
chelates, i.e. the metal ions are complexed,
typically by organic ligands. Examples of these
organic ligands are linear and macrocyclic poly-
aminopolycarboxylic acids and their derivatives.
20 Unfortunately, in many cases there are also
toxicity problems with the free organic ligand.
Thus, even the use of metal chelates as contrast
agents may cause toxicity problems to the extent
that free metal and/or free organic ligand may both
25 be present in the blood following introduction of
the chelate.

 The European patent 0270483 discloses the
use of a formulation excipient of the formula

30




which is described as the metal (M = e.g. sodium)
 salt of a less toxic metal (M' = e.g. zinc, copper
 or calcium) chelate, wherein L may or may not be
 the organic ligand with which the paramagnetic or
 5 heavy metal is complexed. This excipient is
 disclosed as a scavenger for the free metal ions,
 but no mention is made of scavenging free organic
 ligand. Indeed, this European patent suggests and
 claims the addition of free ligand to enhance
 10 safety.

In WO 89/00052 entitled "Metal-Ligand
 Chelates Safety Enhancement - Used in Magnetic
 Resonance Imaging or X-ray Contrast Agents, by
 Addition of Calcium Ions", it is claimed that the
 15 use of effective amounts of calcium in the form of,
 calcium chloride, calcium gluconate, or balanced
 salt solutions substantially reduces the toxicity
 without the need to add additional ligand. The
 potential toxicity from free metal ion and/or
 20 organic ligand was not discussed.

In accordance with the present invention
 novel excipients for metal chelate contrast agents,
 M(L), wherein M is a contrast metal and L is a
 25 chelating ligand, for magnetic resonance, x-ray,
 ultrasound and radiodiagnostic imaging, and
 compositions and methods utilizing such excipients,
 are disclosed. These novel excipients have the
 formula


30



wherein X and X' are each independently calcium or
 zinc;

L' is an organic ligand which may be the organic ligand, L, employed in the metal chelate contrast agent or another organic ligand which has a greater affinity for the metal, M, than for calcium or zinc; and wherein m and n are independently selected from 1, 2 or 3. This salt of the complex of the organic ligand is a highly useful excipient for metal chelate contrast agents in that this single excipient has been found to scavenge free metal ions and free organic ligand, thereby enhancing the safety of such contrast agents and methods employing same.


The present invention pertains to novel excipients for metal chelate contrast agents, compositions of contrast agents complexed with such excipients and methods of imaging employing same. Unexpectedly, the novel excipient of the present invention, comprising the salt of the complex of an organic ligand, has been found to scavenge free metal and free organic ligand. Preferably, the excipient comprises the calcium salt of the calcium complex of the ligand shown as



wherein m, n and L' are as defined above. Since the excipients of the present invention are dual-functioning scavengers and are much safer than the free metal ions and free ligands they scavenge, significantly less toxic contrast agents and methods of imaging are provided.

This dual-functioning phenomenon is additionally advantageous in that the possible toxicity resulting from any dissociation of a metal chelate contrast agent, M(L), which may occur while
5 in storage prior to use is also alleviated. Thus, products with enhanced safety and shelf-life are provided by use of the present excipients.

In viewing the dual scavenging activity of the present excipient
10



it is believed that the calcium or zinc complex, X'(L'), reacts with the toxic metal ion of the
15 contrast agent by a metal ion exchange process. The free calcium or zinc ion of the excipient forms a complex with free organic ligand that may be present. No other new species are expected to form *in situ* utilizing this excipient. The calcium or
20 zinc complex salts of formula I are readily prepared by reacting the desired organic ligand in solution with an excess of a calcium salt, e.g., calcium carbonate, calcium chloride, calcium acetate, zinc chloride, zinc acetate and the like.

25 Thereafter, the present excipients can be employed in compositions with any metal chelate contrast agent comprising a metal ion and an organic ligand. This can be accomplished, using known techniques, by adding the calcium or zinc
30 complex salt of formula I to a solution of the metal chelate contrast agent, as more clearly illustrated in the Examples.


As would be understood by those working in this art, the organic ligand, L', should be selected so that the complexes it forms, i.e., Ca(L'), Zn(L') and/or M(L') are well tolerated.

5 Suitable organic ligands include but are not limited to linear or macrocyclic polyaminopolycarboxylic acids and derivatives thereof. One preferred group metal chelate contrast agents and imaging methods of the present invention employ

10 organic ligands which are 1-substituted-1,4,7-triscarboxymethyl-1,4,7,10-tetraazacyclododecane and derivatives thereof, as disclosed in U. S. 4,885,363 and pending applications U. S. Ser. No. 454,883, filed December 22, 1989 entitled



15 "10-(2'-Hydroxy-3'-Alkoxy-1,4,7-Triscarboxymethyl-1,4,7,10-Tetraazacyclododecanes" and U. S. Ser. No. 454,890, filed December 22, 1989 entitled 10-(2'-Hydroxy-3'-Polyoxaalkyl)-1,4,7-Triscarboxymethyl-1,4,7,10-Tetraazacyclododecane",

20 incorporated herein by reference, which have the general formula





where Y is oxygen or $\overset{\text{R}_1}{\underset{|}{\text{N}}}$;

R₁ and R₂ are each independently hydrogen,
alkyl, arylalkyl, aryl, alkoxy, hydroxyalkyl,
5 hydroxyalkoxy



15 wherein G is NH₂, NCS, $\text{N} - \overset{\text{O}}{\parallel} \text{C} - \underset{\text{H}}{\text{CH}_2} - \text{X}, \text{CO}_2\text{H}, \text{NHR}_4,$
 $\text{N}(\text{R}_4)_2, \text{CN},$ wherein R₄ is alkyl or hydroxyalkyl,




wherein n and m are zero or an integer from one to
five, R₃ is hydrogen, hydroxyalkyl, alkoxy, alkyl,
30 aryl, arylalkyl or hydroxyalkoxy and X is chloro,
bromo or iodo.

R₁ and R₂ are hydrogen in a preferred embodiment for forming a Gd(III) chelate useful in general purpose magnetic resonance imaging. The most preferred embodiment for forming a Gd(III) chelate is when R₁ is hydroxyalkyl or when R₁ is

$$(\text{CH}_2)_n - \overset{\text{O}}{\overset{\parallel}{\text{C}}} (\text{CH}_2)_m - \text{G},$$
 wherein n is 1, m is 0, G is NHR₄ wherein R₄ is alkyl. Thus, preferred ligands from this group are 1,4,7,10-tetraazacyclododecane-1,4,7-triacetic acid, i.e., DO3A, 1,4,7-tris-(carboxymethyl)-10-(2'-hydroxypropyl)-1,4,7,10-tetraazacyclododecane, i.e., HP-DO3A, and 1,4,7-tris(carboxymethyl)-1,4,7,10-tetraazacyclododecane.

Another group of suitable ligands are disclosed in U. S. 4,647,447 which describes the complex salts



or



wherein

X is -COOY, PO₃HY or -CONHOY;

Y is a hydrogen atom, a metal ion equivalent and/or a physiologically biocompatible cation of an inorganic or organic base or amino acid;

A is $-\text{CHR}_2-\text{CHR}_3-$, $-\text{CH}_2\text{CH}_2(\text{ZCH}_2-\text{CH}_2)_m-$,
 $-\text{CH}_2-\overset{\text{N}(\text{CH}_2\text{X})_2}{\underset{|}{\text{CH}}}-\text{CH}_2$, or $-\text{CH}_2-\text{CH}_2-\overset{\text{CH}_2-\text{CH}_2-\text{N}(\text{CH}_2\text{X})_2}{\underset{|}{\text{N}}}-\text{CH}_2-\text{CH}_2-$, wherein
 X is as defined above;


5 each R_1 is hydrogen or methyl;

R_2 and R_3 together represent a trimethylene group or a tetramethylene group or individually are hydrogen atoms, lower alkyl groups (e.g., 1-8 carbons), phenyl groups, benzyl groups or R_2 is a
 10 hydrogen atom and R_3 is $-(\text{CH}_2)_p-\text{C}_6\text{H}_4-\text{W}-\text{protein}$ where p is 0 or 1, W is $-\text{NH}-$, $-\text{NHCOCH}_2-$ or $-\text{NHCS}-$, protein represents a protein residue;

m is 1, 2 or 3;

15 Z is an oxygen atom or a sulfur atom or the group NCH_2X or $\text{NCH}_2\text{CH}_2\text{OR}_4$ wherein X is as defined above and R_4 is C_{1-8} alkyl;

V is X or is $-\text{CH}_2\text{OH}$, $-\text{CONH}(\text{CH}_2)_n\text{X}$ or $-\text{COB}$, wherein X is as defined above, B is a protein or lipid residue, n is an integer from 1 to 12, or if
 20 R_1 , R_2 and R_3 are each hydrogen; then both V's together form the group




25

where X is as above, w is 1, 2 or 3, provided that at least two of the substituents Y represent metal ion equivalents of an element with an atomic number of 21 to 29, 42, 44 or 57 to 83. Preferred ligands
 30 from U. S. 4,647,447 include 1,4,7,10-tetraazacyclododecane- $\text{N},\text{N}',\text{N}'',\text{N}'''$ -tetraacetic acid, i.e., DOTA, and diethylene triamine pentaacetic acid, i.e., DTPA.

Also related to the U. S. 4,647,447 ligands are diethylenetriamine pentaacetic acid-bis methanamide (DTPA-BMA), which can be shown as


5



10

DTPA-bis morpholino amide ((a) below); and DTPA-bis 1,2-dihydroxypropylamide ((b) below)

15




20

a) R_1 and R_2 together with nitrogen to which they are attached form morpholino,


b) R_1 is hydrogen and R_2 is

25



Another group of ligands suitable for use with the present excipients/compositions/methods is disclosed in European Application 0 255 471 A1. That application discloses macrocycles of the formula

30



Y is N or P;

A^1 and A^2 are each optionally branched C_{2-6}

U^1, U^2, U^3 and U^4 are each a single bond or

D¹, D², D³, D⁴ are each O, S, C₁₋₆ alkylene

R₇ is hydrogen or C₁₋₄ alkylene having a

R¹ is hydrogen or a metal ion equivalent;

D⁵ is D¹ or CHR⁵, where R⁵ can be hydrogen or optionally unsaturated C₁₋₂₀ alkylene which may be imino, phenyleneoxy, phenyleneimino, amido, O, S and/or N optionally substituted with imino and/or amino and may carry a terminal functional group (optionally bonded to a

s and t are each 0-5;

R₂ is hydrogen, optionally substituted C₁₋₁₆ acyl, acylalkyl (optionally substituted by more OH or lower alkoxy groups), -CH₂-X-V, BOB where X is CO, optionally branched C₁₋₁₀ (optionally substituted by 1 or more OH or alkoxy groups) or optionally branched C₂₋₂₃


V is NR^3R^4 or COOR^6 ;

R³ and R⁴ are each hydrogen, C₁₋₁₆ alkyl (optionally substituted by 1 or more OH or lower alkoxy groups) or together complete a 5-6 membered heterocycle optionally containing another
 5 heteroatom;

R₆ is hydrogen, C₁₋₁₆ saturated, unsaturated, linear branched or cyclic hydrocarbyl, aryl or aralkyl;

R₂ or R₃ can be bonded by a C₂₋₂₀ alkylene chain (optionally having a terminal carbonyl group, optionally interrupted by 1 or more O or R¹ carboxymethylimino, or substituted by one or more OH, lower alkoxy or carboxy lower alkyl groups) to
 10 a second macromolecule of the formula

15



20 which second macromolecule D' can be the same as or different from the macromolecule of D.

As described above, the excipients of the present invention can be employed with any metal chelate contrast agent. In one embodiment they are
 25 used with a contrast agent comprising an organic ligand of formula A complexed with a paramagnetic metal atom and used as relaxation enhancement agents for magnetic resonance imaging. These agents, when administered to a mammalian host
 30 (e.g., humans) distribute in various concentrations to different tissues, and catalyze relaxation of protons (in the tissues) that have been excited by

the absorption of radiofrequency energy from a magnetic resonance imager. This acceleration of the rate of relaxation of the excited protons provides for an image of different contrast when
5 the host is scanned with a magnetic resonance imager. The magnetic resonance imager is used to record images at various times generally before and after administration of the agents, and the differences in the images created by the agents'
10 presence in tissues are used in diagnosis. In proton magnetic resonance imaging, paramagnetic metal atoms such as gadolinium(III), dysprosium(III), manganese(II), manganese(III) chromium(III), iron(II) and iron(III) (all are paramagnetic metal
15 atoms with a symmetrical electronic configuration) are preferred as metals complexed by the ligands of formula I; gadolinium(III) is most preferred due to the fact that it has the highest paramagnetism, low toxicity, and high lability of coordinated water.

20 Exemplary contrast agents which will be greatly enhanced when employed in a composition including a pharmaceutically acceptable carrier and an excipient of the present invention include Gadoteridol which is $\text{Gd}(\text{HP-DO3A})$, Dotarem which is
25 N-methylglucamine $[\text{Gd}(\text{DOTA})]$, Magnevist which is di-N-methylglucamine $[\text{Gd}(\text{DTPA})]$ and Gadodiamide which is $\text{Gd}(\text{DTPA-BMA})$, $\text{Gd}(\text{DTPA})$ -bis morpholino-amide, $\text{Gd}(\text{DTPA})$ -bis 1,2-dihydroxypropylamide.

30 Additionally, the excipients of formula I are conveniently employed with ligands of formula A which are complexed with a lanthanide (atomic

number 58 to 71) and used as chemical shift agents in magnetic resonance imaging or in magnetic resonance *in vivo* spectroscopy.

Excipients of formula I are also
5 conveniently employed with contrast elements including yttrium and the transition series (atomic number 21-29).

While the above-described uses for the excipients of formula I and contrast agents
10 including same are preferred, those working in the medical diagnostic arts will appreciate that the excipients can also be used with contrast agents in x-ray imaging, radionuclide imaging and ultrasound imaging.

15 As mentioned previously excipients, compositions and methods wherein $X = X' = \text{calcium}$ are preferred and these excipients, compositions and methods where $L' = L$ are most preferred.

As described previously, contrast agents
20 which include the excipient of formula I can be easily prepared by adding the calcium or zinc complex salt into the metal chelate contrast agent solution. Preferably the so-formed solution is maintained at about pH neutral. The
25 excipient/contrast agent composition can be prepackaged in combination or the present excipient can be added to the contrast solution directly before use. Typically the mole ratio of the calcium or zinc complex salt to the contrast agent
30 is about 0.05-10 percent.

The following Examples illustrate specific embodiments of the present invention, however, it is understood that the invention should not be limited to the details therein.

Example 1

Calcium bis[1,4,7,10-tetraazacyclododecane-1,4,7-triacetatocalcium(II)], $\text{Ca}[\text{Ca}(\text{DO3A})]_2^*$

- 5 Twenty millimoles (7.67 g) of DO3A was dissolved in 32 mL of water and 32 mmoles (3.18 g) of calcium carbonate was added slowly. The solution was heated under reflux at 80°C for 2 hours and it was cooled to room temperature and filtered to
10 remove excess solid calcium carbonate. The solution was then rotary evaporated to dryness and placed in a 75°C vacuum oven (5 mm Hg) for 24 hours. A glass-like solid was obtained in 94 percent yield.
- 15 Analysis calc'd $\text{Ca}_3\text{C}_{28}\text{H}_{46}\text{N}_8\text{O}_{12}$ (8% H_2O):
 C, 38.58; H, 6.15; N, 12.85;
Found: C, 38.17; H, 5.99; N, 12.83.

Example 2

20

Calcium bis[1,4,7-tris(carboxymethyl)-10-(2'-hydroxy)propyl-1,4,7,10-tetraazacyclododecanatocalcium(II)], $\text{Ca}[\text{Ca}(\text{HP-DO3A})]_2^{**}$

- The ligand HP-DO3A, containing 3.5 percent
25 water by elemental analysis data, (0.518 g, 1.24 mmol) was dissolved in 10 mL water at room temperature and solid calcium carbonate (0.204 g, 2.03 mmol) was added to it slowly with stirring. The cloudy solution was heated at 90°C for 2.5
30 hours, cooled, centrifuged, and filtered. The filtrate was evaporated under vacuum to dryness to give 0.56 g of white solid. The solid was recrystallized in water:acetone (1:3 v/v) mixture and dried *in vacuo* at room temperature.

Analysis calc'd $\text{Ca}_3\text{C}_{34}\text{H}_{58}\text{N}_8\text{O}_{14}$ (12% H_2O):

C, 39.50; H, 6.85; N, 10.84;

Found: C, 39.66; H, 6.83; N, 10.76.

5 *DO3A = 1,4,7,10-tetraazacyclododecane-1,4,7-
 triacetate

10 **HP-DO3A = 1,4,7-tris(carboxymethyl)-10-(2'-
 hydroxy)propyl-1,4,7,10-tetraazacyclo-
 dodecanoate.

Example 3

15 Calcium[1,4,7,10-tetraazacyclododecane-1,4,7-10-
 tetraacetatocalcium(II)], $\text{Ca}[\text{Ca}(\text{DOTA})]^*$

20 Twenty millimoles of DOTA is dissolved in 40
 mL of water and 44 mmoles of CaCO_3 is added slowly.
 The solution is heated under reflux at 80°C for 2
 hours and it is cooled to room temperature and
 filtered to remove excess solid CaCO_3 . The solution
 is then rotary evaporated to dryness and placed in
 a 75°C vacuum oven (5 mm Hg) for 24 hours. A white
 solid is obtained.

25 *DOTA = 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-
 tetraacetate

Example 4

Triscalcium bis[N,N'-bis[2-[bis(carboxymethyl)-
amino]ethylglycinatocalcium(II)], $\text{Ca}_3[\text{Ca}(\text{DTPA})]_2$ **

5

Twenty millimoles of DTPA is dissolved in 40 mL of water and 55 mmoles of CaCO_3 is added slowly. The solution is heated under reflux at 80°C for 2 hours and it is cooled to room temperature and
10 filtered to remove excess solid CaCO_3 . The solution is then rotary evaporated to dryness and placed in a 75°C vacuum oven (5 mm Hg) for 24 hours. A white solid is obtained.

15 **DTPA = N,N-bis[2-[bis(carboxymethyl)-amino]ethylglycinate

Example 5

The relative toxicities of free metal, free
20 ligand, metal chelate contrast agent and excipient of the present invention were determined by injecting samples of each into the tail vein of Charles River CD-1 mice (18-23 g, 25-35 days) at 0.02 mL/g. The observation period was 14 days.

25

The results are summarized in Table 1.

Table 1

Intravenous Acute Tolerance in Mice (LD₅₀ Values)
 for Gd(OH)₃, DO3A, DOTA, Gd(DO3A),
 5 Gd(DO3A) formulated, Gd(HP-DO3A),
 Gd(HP-DO3A) formulated, Ca[Ca(DO3A)]₂,
Ca[Ca(HP-DO3A)]₂, and CaCl₂ at the Physiologic pH.

	Compound	LD ₅₀ (mMol/kg)
10	Gd(OH) ₃	0.1
	DO3A	0.12
	HP-DO3A	0.11
	DOTA	0.18
15	Gd(DO3A)	5.7
	Gd(DO3A), formulated ^a	7.4 (male)
		8.5 (female)
	Gd(HP-DO3A)	10
	Gd(HP-DO3A), formulated ^b	10.7 (male)
20		13.6 (female)
	Ca[Ca(DO3A)] ₂	1.6
	Ca[Ca(HP-DO3A)] ₂	1.3
	CaCl ₂	1.5

^a The formulation consists of Gd(DO3A), 0.5 M;
 25 Ca[Ca(DO3A)]₂, 0.25 mM; Tris Buffer, 10 mM (pH
 7.4).

^b The formulation consists of Gd(HP-DO3A), 0.5 M;
 Ca[Ca(HP-DO3A)]₂, 0.25 mM; Tris Buffer, 10 mM (pH
 7.4).

The data illustrate that the complexed Gd(DO3A) (i.e., metal chelate contrast agent) and calcium complex salt (i.e., excipient) were highly tolerated. Therefore, when using the calcium
5 complex salt of the present invention as the excipient, the amounts of free metal and free ligand are greatly reduced and any excesses of chelated contrast agent or excipient are well tolerated providing contrast methodology with
10 enhanced safety.

Example 6

The following novel diagnostic formulation
15 was prepared by combining the known diagnostic agent, Gadoteridol, with the excipient of Example 2 using the procedure which follows:

	<u>Ingredient</u>	<u>Amount (per ml)</u>
20	Gadoteridol ¹ (diagnostic agent)	558.6 mg
	Calteridol Calcium ² (novel excipient)	0.46 mg
	Tromethamine (buffer)	1.21 mg
25	1N NaOH solution	as needed for pH adjustment
	1N HCl solution	as needed for pH adjustment
	Water for Injection	1.0 ml
30	USP q.s.	

¹Gadolinium(III) 1,4,7-tris(carboxymethyl)-10-(2'-hydroxypropyl)-1,4,7,10-tetraazacyclododecane

²Ca[Ca(HP-DO3A)]₂ excipient of Example 2 above.

Procedure: About 60% of the necessary water was heated to between 48 and 52°C and vigorously agitated while adding the buffer (tromethamine) and excipient (calteridol calcium). Thereafter, the
5 diagnostic agent (gadoteridol) was added slowly and the agitation was continued for about 1 hour while maintaining the temperature at 48-52°C. The so-prepared solution was cooled to 20-25°C and the pH was adjusted to between 7.3 and 7.5 (7.4
10 optimum) using HCl solution and/or NaOH solution as necessary. The balance of the water was added and the pH rechecked/readjusted. The solution was filtered through a sterile 0.2 micrometer membrane and thereafter sterilized and stored at controlled
15 room temperature (15-30°C).

Example 7

A novel diagnostic composition for Dotarem,
20 i.e., N-methylglucamine [Gd(DOTA)], was prepared using the procedure above in Example 6 but substituting Dotarem for Gadoteridol and substituting the excipient of Example 3, Ca[Ca(DOTA)], for the calteridol calcium.

25

Example 8

A novel diagnostic composition for Magnevist, i.e., (N-methylglucamine)₂ [Gd(DTPA)],
30 was prepared using the procedure of Example 6 but substituting Magnevist for Gadoteridol and substituting the excipient of Example 4, Ca₃[Ca(DTPA)]₂, for the calteridol calcium.

Example 9

A novel diagnostic composition for Gadodiamide, i.e., $\text{Gd}(\text{DTPA-bis methylamide})$, was prepared using the procedure of Example 6 but substituting Gadodiamide for Gadoteridol and substituting an excipient $\text{Ca}[\text{Ca}(\text{DTPA-bis methylamide})]_2$, prepared using the methodology of Example 2.

10

Example 10

A novel diagnostic composition was prepared for the contrast agent gadolinium DTPA bis morpholinoamide using the procedure of Example 6 substituting this agent for Gadoteridol and substituting an excipient, $\text{Ca}[\text{Ca}(\text{DTPA bis morpholinoamide})]_2$ prepared using the methodology of Example 2.

15

Example 11

A novel diagnostic composition was prepared for the contrast agent gadolinium (DTPA 1,2-di-hydroxypropylamide) using the procedure of Example 6 substituting this agent for Gadoteridol and substituting an excipient, $\text{Ca}[\text{Ca}(\text{DTPA 1,2-di-hydroxypropylamide})]_2$, prepared using the methodology of Example 2.

20

25